

LAMPIRAN



PAK LI
MAL13115185TC
4 Sheets
Size: 10 cm x 7 cm

Digunakan secara tradisional untuk melegakan sakit otot, sakit bahu, sakit belakang dan sakit sendi.

Pak Li Medicated Plaster

李太医膏药贴

傳統上用於緩解肌肉酸痛，肩膊酸痛，背痛和關節疼痛。

特点 / Ciri-ciri:

- ✓ 伸縮自如 *Mudahalih*
- ✓ 彈性更佳 *Lebih fleksibel*
- ✓ 柔軟舒適 *Lebih selesa*
- ✓ 易撕易貼 *Mudah ditanggal & dilekatkan*
- ✓ 黏性特佳 *Lebih melekat*

VALUE PACK

Indikasi: Digunakan secara tradisional untuk melegakan sakit otot, sakit bahu, sakit belakang dan sakit sendi.

Cara Penggunaan: Bersih dan keringkan tempat kesakitan, tanggalkan lapisan perlindungan dan lekat pada tempat sakit. Sila gunakan produk ini satu atau dua kali sehari.

Perhatian: Untuk kegunaan luar sahaja. Jauh daripada kanak-kanak. Sila merujuk kepada ahli farmasi / doktor jika simptom berlarutan / bertambah teruk. Simpan di tempat di bawah 30°C. Lindunginya daripada haba dan kelembapan.

《功效》傳統上用於緩解肌肉酸痛，肩膊酸痛，背痛和關節疼痛。
《使用方法》用前將不適處洗淨擦干，揭去藥貼覆膜，粘貼於不適之處，用手按牢即可。
《注意事項》請將本品放在兒童不能接觸到的地方，本品屬外用產品，禁止內服。如病情持續，請即諮詢醫師，置於陰涼乾燥處。

Indication: Traditionally used for relieving muscular ache, shoulder ache, backache and joint pain.

Administration: Clean and dry the affected area properly. Remove the paper attached to the plaster and apply. Please use the product once or twice daily.

Precaution: For external used only. This is traditional medicine. Please consult your pharmacist / doctor if symptoms persist / worsen. Keep out of reach of children. Store in areas below 30°C. Protect from light and moisture.

Contra-indication: Sensitivity to any of the plaster ingredients.

Pregnancy and lactation: Not recommended due to lack of sufficient data.

WARNING: CAN CAUSE CONVULSION. CONTRAINDICATED IN INFANTS BELOW 2 YEARS OF AGE. CAUTION MUST BE EXERCISED WHEN OLDER CHILDREN ARE TREATED. AVOID DIRECT APPLICATION INTO NOSTRILS.
It is dangerous to place any camphor-containing product into the nostril of children. A small amount applied this way may cause immediate collapse. Avoid contact with the eyes. Do not apply to wounds or damaged skin.

Ingredient (per cm²) contains:

Sanguis Draxonis	0.66mg
Radix Notoginseng	0.22mg

Batch No.: C-21810601
Mfg. Date: JUN 2018
Exp. Date: JUN 2021

Manufactured by:
PT Indo Abadi Sarmasmar, Dusun Ngabih, Desa Gempolkuring, Kec. Menganti, Kabupaten Gresik, Indonesia.
Product license holder and distributor:
Tong Huat Medical Sdn. Bhd.
4748-4741, Jalan Telaga Air, 12200 Butterworth, Pulau Pinang, Malaysia.

李太医膏药贴

Pak Li Medicated Plaster



PAK LI
MAL13115185TC
4 Sheets
Size: 10 cm x 7 cm

Digunakan secara tradisional untuk melegakan sakit otot, sakit bahu, sakit belakang dan sakit sendi.

Pak Li Medicated Plaster

李太医膏药贴

傳統上用於緩解肌肉酸痛，肩膊酸痛，背痛和關節疼痛。

特点 / Ciri-ciri:

- ✓ 伸縮自如 *Mudahalih*
- ✓ 彈性更佳 *Lebih fleksibel*
- ✓ 柔軟舒適 *Lebih selesa*
- ✓ 易撕易貼 *Mudah ditanggal & dilekatkan*
- ✓ 黏性特佳 *Lebih melekat*

Indikasi: Digunakan secara tradisional untuk melegakan sakit otot, sakit bahu, sakit belakang dan sakit sendi.

Cara Penggunaan: Bersih dan keringkan tempat kesakitan, tanggalkan lapisan perlindungan dan lekat pada tempat sakit. Sila gunakan produk ini satu atau dua kali sehari.

Perhatian: Untuk kegunaan luar sahaja. Jauh daripada kanak-kanak. Sila merujuk kepada ahli farmasi / doktor jika simptom berlarutan / bertambah teruk. Simpan di tempat di bawah 30°C. Lindunginya daripada haba dan kelembapan.

《功效》傳統上用於緩解肌肉酸痛，肩膊酸痛，背痛和關節疼痛。
《使用方法》用前將不適處洗淨擦干，揭去藥貼覆膜，粘貼於不適之處，用手按牢即可。
《注意事項》請將本品放在兒童不能接觸到的地方，本品屬外用產品，禁止內服。如病情持續，請即諮詢醫師，置於陰涼乾燥處。

Indication: Traditionally used for relieving muscular ache, shoulder ache, backache and joint pain.

Administration: Clean and dry the affected area properly. Remove the paper attached to the plaster and apply. Please use the product once or twice daily.

Precaution: For external used only. This is traditional medicine. Please consult your pharmacist / doctor if symptoms persist / worsen. Keep out of reach of children. Store in areas below 30°C. Protect from light and moisture.

Contra-indication: Sensitivity to any of the plaster ingredients.

Pregnancy and lactation: Not recommended due to lack of sufficient data.

WARNING: CAN CAUSE CONVULSION. CONTRAINDICATED IN INFANTS BELOW 2 YEARS OF AGE. CAUTION MUST BE EXERCISED WHEN OLDER CHILDREN ARE TREATED. AVOID DIRECT APPLICATION INTO NOSTRILS.
It is dangerous to place any camphor-containing product into the nostril of children. A small amount applied this way may cause immediate collapse. Avoid contact with the eyes. Do not apply to wounds or damaged skin.

Ingredient (per cm²) contains:

Sanguis Draxonis	0.66mg
Radix Notoginseng	0.22mg

Batch No.: C-21810601
Mfg. Date: JUN 2018
Exp. Date: JUN 2021

Manufactured by:
PT Indo Abadi Sarmasmar, Dusun Ngabih, Desa Gempolkuring, Kec. Menganti, Kabupaten Gresik, Indonesia.
Product license holder and distributor:
Tong Huat Medical Sdn. Bhd.
4748-4741, Jalan Telaga Air, 12200 Butterworth, Pulau Pinang, Malaysia.